

GUJARAT TECHNOLOGICAL UNIVERSITY

BE - SEMESTER- 1st / 2nd EXAMINATION (NEW SYLLABUS) – SUMMER 2018

Subject Code: 2110015

Date: 17-05-2018

Subject Name: Vector Calculus and Linear Algebra

Time: 02:30 pm to 05:30 pm

Total Marks: 70

Instructions:

1. Question No. 1 is compulsory. Attempt any four out of remaining Six questions.
2. Make suitable assumptions wherever necessary.
3. Figures to the right indicate full marks.

Q.1	Objective Question (MCQ)	Mark
------------	---------------------------------	-------------

(a)	Choose the appropriate answer for the following questions.	07
------------	--	-----------

1. A square matrix whose determinant is non zero is called
(A) Singular (B) non-singular (C) invertible (D) both B and C
2. If A and B are non singular matrices then $(AB)^{-1} = \text{_____}$
(A) $A^{-1}B^{-1}$ (B) AB (C) $B^{-1}A^{-1}$ (D) none of these
3. If $A = \begin{bmatrix} 1 & 0 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{bmatrix}$ then A is in
(A) Row echelon form (B) Reduced Row echelon form (C) both A and B (D) none of these
4. For what values of k does the system $x + y = 2, 3x + 3y = k$ has infinitely many solutions
(A) K=5 (B) k=4 (C) k=6 (D) k=1
5. If in a set of vectors atleast one member can be expressed as a linear combination of the remaining vectors then the set is
(A) Linearly independent (B) Linearly dependent (C) basis (D) none of these
6. If V is any vector space and S be a subset of V then S is called basis for V if
(A) S is Linearly independent (B) S spans V (C) both A and B
(D) S is Linearly dependent
7. For what value of k the vectors u and v are orthogonal where $u=(2,1,3), v=(1,7, k)$
(A) K=-3 (B) k=1 (C) k=5 (D) k=2

(b)	Choose the appropriate answer for the following questions.	07
------------	--	-----------

1. The eigen values of a matrix $A = \begin{bmatrix} 1 & 0 \\ 2 & 4 \end{bmatrix}$ are
(A) 1,4 (B) -1,-4 (C) 1,3 (D) -1,3
2. If A is a nxn size invertible matrix then rank of A is
(A) n-1 (B) n (C) 2n (D) n+1

3. If \overline{F} is solenoidal then
(A) $\nabla \overline{F} = 0$ (B) $\nabla \times \overline{F} = 0$ (C) $\nabla \cdot \overline{F} = 0$ (D) none of these
4. The mapping $T : R^3 \rightarrow R^3$ defined by $T(x, y, z) = (x, y, -z)$ is called as
(A) Contraction (B) Projection (C) Reflection (D) Rotation
5. The linear transformation $T : V \rightarrow W$ is one to one if and only if the nullspace of T consists of only
(A) Identity vector (B) zero vector (C) any non zero vector (D) none of these
6. If $A = \begin{bmatrix} 1 & 1 \\ 2 & 2 \end{bmatrix}$ then the rank of the matrix A is
(A) 1 (B) 2 (C) 0 (D) 4
7. Let A be a skew-symmetric matrix then
(A) $a_{ij} = a_{ji}$ (B) $a_{ij} = -a_{ji}$ (C) $a_{ii} = 0$ (D) both B and C

Q.2 (a) Find the unit vector normal to the surface $xy^3z^2 = 4$ at $(-1, -1, 2)$ **03**

(b) Express the matrix $A = \begin{bmatrix} 3 & -2 & 6 \\ 2 & 7 & -1 \\ 5 & 4 & 0 \end{bmatrix}$ as the sum of a symmetric and skew-symmetric matrix. **04**

(c) Investigate for what values of λ and μ the equations **07**
 $2x + 3y + 5z = 9, 7x + 3y - 2z = 8, 2x + 3y + \lambda z = \mu$ have
(1) No solution (2) a unique solution (3) infinite number of solutions

Q.3 (a) Find the rank of the matrix $\begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{bmatrix}$ **03**

(b) Find the inverse of the matrix $\begin{bmatrix} 2 & 3 & 4 \\ 4 & 3 & 1 \\ 1 & 2 & 4 \end{bmatrix}$ by Gauss Jordan Method **04**

(c) For the basis $S = \{v_1, v_2, v_3\}$ of R^3 where **07**
 $v_1 = (1, 1, 1), v_2 = (1, 1, 0), v_3 = (1, 0, 0)$ Let $T : R^3 \rightarrow R^2$ be the linear transformation such that
 $T(v_1) = (1, 0), T(v_2) = (2, -1), T(v_3) = (4, 3)$ find a formula for $T(x_1, x_2, x_3)$ and then use the formula to find $T(4, 3, -2)$

- Q.4 (a)** Determine whether the vector $v = (-5, 1, -7)$ is a linear combination of the vectors $v_1 = (1, -2, 2), v_2 = (0, 5, 5), v_3 = (2, 0, 8)$ **03**
- (b)** Solve the linear system $x + y + z = 4, -x - y + z = -2, 2x - y + 2z = 2$ by gauss elimination method. **04**
- (c)** Let R^3 have the Euclidean inner product. Use the gram schmidt process to transform the basis (u_1, u_2, u_3) in to Orthonormal basis where $u_1 = (1, 0, 0), u_2 = (3, 7, -2), u_3 = (0, 4, 1)$ **07**
- Q.5 (a)** Find the eigen values and corresponding eigen vectors of $A = \begin{bmatrix} 2 & -12 \\ 1 & -5 \end{bmatrix}$ **03**
- (b)** Let $A = \begin{bmatrix} -2 & 3 \\ 1 & -2 \\ 1 & -1 \end{bmatrix}$ and $b = \begin{bmatrix} 1 \\ -1 \\ 2 \end{bmatrix}$ then find the least squares solutions to $AX=b$ **04**
- (c)** Let $T : R^3 \rightarrow R^3$ be a linear operator and $B = (v_1, v_2, v_3)$ a basis for R^3 . Suppose that $T(v_1) = (1, 1, 0), T(v_2) = (1, 0, -1), T(v_3) = (2, 1, -1)$ then **(1) Is $(1, 2, 1)$ in $R(T)$? (2) Find a basis for $R(T)$.** **07**
- Q.6 (a)** Find the work done by the force $\vec{F} = (3x^2 - 3x)i + 3zj + k$ along the straight line $ti + tj + tk, 0 \leq t \leq 1$. **03**
- (b)** Check whether the vectors $(2, -3, 1), (4, 1, 1), (0, -7, 1)$ is a basis for R^3 **04**
- (c) Verify Green's Theorem for** $\vec{F} = (x - y)i + xj$ and C is $x^2 + y^2 = 1$ **07**
- Q.7 (a)** Find the directional derivative of $4xz^2 + x^2yz$ at $(1, -2, -1)$ in the direction of $2i - j - 2k$ **03**
- (b)** Show that $\vec{F} = (e^x \cos y + yz)i + (xz - e^x \sin y)j + (xy + z)k$ is conservative and find the potential function. **04**
- (c)** Let $V = \{(a, b) / a, b \in R\}$ and let $v = (v_1, v_2), w = (w_1, w_2)$ then define $(v_1, v_2) + (w_1, w_2) = (v_1 + w_1 + 1, v_2 + w_2 + 1)$ and $c(v_1, v_2) = (cv_1 + c - 1, cv_2 + c - 1)$ then verify that V is a vector space. **07**
